

Clinical Article: August 15, 2008

A decade with Silicone Hydrogels Part 1

References

1. Sweeney D, du Toit R, Keay L et al. *Clinical performance of silicone hydrogel lenses*. in *Silicone hydrogels: Continuous wear contact lenses*, D. Sweeney, Editor. Oxford, Butterworth-Heinemann,2004, pp 164 - 216.
2. Stapleton F, Stretton S, Papas E et al. Silicone hydrogel contact lenses and the ocular surface. *Ocul Surf* 2006; 4;1: 24-43.
3. Morgan P. Trends in UK contact lens prescribing 2007. *Optician* 2007; 233 (6104): 16-17
4. Morgan P. Trends in UK contact lens prescribing 2008. *Optician* 2008; 235 (6155): 18-19
5. Poggio EC, Glynn RJ, Schein Od et al. The incidence of ulcerative keratitis among users of daily-wear and extended-wear soft contact lenses. *N Engl J Med* 1989; 321;12: 779-83.
6. Schein OD, Glynn RJ, Poggio EC et al. The relative risk of ulcerative keratitis among users of daily-wear and extended-wear soft contact lenses. A case-control study. *Microbial Keratitis Study Group. N Engl J Med* 1989; 321;12: 773-8.
7. Cheng KH, Leung SL, Hoekman HW et al. Incidence of contact-lens-associated microbial keratitis and its related morbidity. *Lancet* 1999; 354;9174: 181-5.
8. Dart JK, Radford CF, Minassian D et al. Risk Factors for Microbial Keratitis with Contemporary Contact Lenses A Case-Control Study. *Ophthalmology* 2008.
9. Stapleton F, Keay L, Edwards K et al. The Incidence of Contact Lens-Related Microbial Keratitis in Australia. *Ophthalmology* 2008.
10. Tighe B: *Contact Lens Materials*. in *Contact Lenses*, A. Phillips and L. Speedwell, Editors. Edinburgh, Butterworth-Heinemann,2006, pp 59 - 78.
11. Tighe B: *Silicone hydrogels: Structure, properties and behaviour*. in *Silicone Hydrogels: Continuous Wear Contact Lenses*, D. Sweeney, Editor. Oxford, Butterworth-Heinemann,2004, pp 1 - 27.
12. Efron N, Morgan PB, Cameron ID et al. Oxygen permeability and water content of silicone hydrogel contact lens materials. *Optom Vis Sci* 2007; 84;4: 328-37.
13. Holden BA, Mertz GW. Critical oxygen levels to avoid corneal edema for daily and extended wear contact lenses. *Invest Ophthalmol Vis Sci*. 1984; 25 (10): 1161-7
14. Fonn D, Bruce AS: A review of the Holden-Mertz criteria for critical oxygen transmission. *Eye Contact Lens* 2005; 31;6: 247-51.
15. Brennan NA: Beyond flux: total corneal oxygen consumption as an index of corneal oxygenation during contact lens wear. *Optom Vis Sci* 2005; 82;6: 467-72.
16. Brennan NA: Corneal oxygenation during contact lens wear: comparison of diffusion and EOP-based flux models. *Clin Exp Optom* 2005; 88;2: 103-8.
17. Morgan P, Brennan N: The decay of Dk ? *Optician* 2004; 227;5937: 27 - 33.
18. Efron, N. and Carney, L.G. Oxygen levels beneath the closed eyelid. *Invest Ophthalmol Vis Sci*

1979; 18: 93-95

19. Fonn D, du Toit R, Simpson TL et al. Sympathetic swelling response of the control eye to soft lenses in the other eye. *Invest Ophthalmol Vis Sci* 1999; 40;13: 3116 - 3121.
20. Fonn D, MacDonald KE, Richter D, Pritchard N: The ocular response to extended wear of a high Dk silicone hydrogel contact lens. *Clin Exp Optom* 2002; 85;3: 176-82.
21. Moezzi AM, Fonn D, Simpson TL: Overnight corneal swelling with silicone hydrogel contact lenses with high oxygen transmissibility. *Eye Contact Lens* 2006; 32;6: 277-80.
22. Steffen RB, Schnider CM: The impact of silicone hydrogel materials on overnight corneal swelling. *Eye Contact Lens* 2007; 33;3: 115-20.
23. Hamano H, Maeda N, Hamano T et al. Corneal thickness change induced by dozing while wearing hydrogel and silicone hydrogel lenses. *Eye Contact Lens* 2008; 34;1: 56-60.
24. Comstock TL, Robboy M, Cox I and Brennan N. Overnight clinical performance of a high Dk silicone hydrogel contact lens. *Contact lens and anterior eye*, 1999; 22 (4): 159
25. Covey M, Sweeney DF, Terry R et al. Hypoxic effects on the anterior eye of high-Dk soft contact lens wearers are negligible. *Optom Vis Sci* 2001; 78;2: 95-99.
26. Keay L, Jalbert I, Sweeney DF, Holden BA: Microcysts: clinical significance and differential diagnosis. *Optometry* 2001; 72;7: 452-60.
27. Stretton S, Jalbert I, Sweeney DF: Corneal hypoxia secondary to contact lenses: the effect of high-Dk lenses. *Ophthalmol Clin North Am* 2003; 16;3: 327-40
28. Fonn D, Sweeney D, Holden BA, Cavanagh D: Corneal oxygen deficiency. *Eye Contact Lens* 2005; 31;1: 23-7.
29. Sweeney DF: Clinical signs of hypoxia with high-Dk soft lens extended wear: is the cornea convinced? *Eye Contact Lens* 2003; 29;1 Suppl: S22-25.
30. Dumbleton KA, Chalmers RL, Richter DB, Fonn D: Vascular response to extended wear of hydrogel lenses with high and low oxygen permeability. *Optom Vis Sci* 2001; 78;3: 147 - 151.
31. Papas EB, Vajdic CM, Austen R, Holden BA: High-oxygen-transmissibility soft contact lenses do not induce limbal hyperaemia. *Curr Eye Res* 1997; 16;9: 942 - 948.
32. Nilsson SE: Seven-day extended wear and 30-day continuous wear of high oxygen transmissibility soft silicone hydrogel contact lenses: a randomized 1-year study of 504 patients. *CLAO J* 2001; 27;3: 125-136.
33. Brennan NA, Coles ML, Connor HR, McIlroy RG: A 12-month prospective clinical trial of comfilcon A silicone-hydrogel contact lenses worn on a 30-day continuous wear basis. *Cont Lens Anterior Eye* 2007; 30;2: 108-18.
34. Fonn D, MacDonald KE, *et al.* The ocular response to extended wear of a high Dk silicone hydrogel contact lens. *Clin Exp Optom* 2002; 85 (3): 176-82
35. Morgan PB, Efron N. Comparative clinical performance of two silicone hydrogel contact lenses for continuous wear. *Clin Exp Optom* 2002; 85 (3): 183-92
36. Papas E, Vajdic C, *et al.* High oxygen-transmissibility soft contact lenses do not induce limbal hyperaemia. *Curr Eye Res* 1997; 16 (9): 942-8

37. Papas E. On the relationship between soft contact lens oxygen transmissibility and induced limbal hyperaemia. *Exp Eye Res* 1998; 67 (2): 125-31
38. du Toit R, Simpson TL, *et al.* Recovery from hyperaemia after overnight wear of low and high transmissibility hydrogel lenses. *Curr Eye Res* 2001; 22 (1): 68-73
39. Riley C, Young G, Chalmers R: Prevalence of ocular surface symptoms, signs, and uncomfortable hours of wear in contact lens wearers: the effect of refitting with daily-wear silicone hydrogel lenses (senofilcon a). *Eye Contact Lens* 2006; 32;6: 281-6.
40. Dumbleton K, Keir N, Moezzi A *et al.* Objective and subjective responses in patients refitted to daily-wear silicone hydrogel contact lenses. *Optom Vis Sci* 2006; 83;10: 758-68.
41. Dillehay SM, Miller MB: Performance of lotrafilcon B silicone hydrogel contact lenses in experienced low-Dk/t daily lens wearers. *Eye Contact Lens* 2007; 33;6 Pt 1: 272-7.
42. Long B, McNally J: The clinical performance of a silicone hydrogel lens for daily wear in an Asian population. *Eye Contact Lens* 2006; 32;2: 65-71.
43. Nilsson SE. Seven-day extended wear and 30-day continuous wear of high oxygen transmissibility soft silicone hydrogel contact lenses: a randomized 1-year study of 504 patients. *CLAO J* 2001; 27 (3): 125-36
44. Covey M, Sweeney DF, *et al.* Hypoxic effects on the anterior eye of high-Dk soft contact lens wearers are negligible. *Optom Vis Sci* 2001; 78 (2): 95-9
45. Maldonado-Codina C, Morgan PB, *et al.* Short-term physiologic response in neophyte subjects fitted with hydrogel and silicone hydrogel contact lenses. *Optom Vis Sci* 2004; 81 (12): 911-21
46. Dumbleton K, Keir N, *et al.* Redness, dryness and comfort following refitting long term low Dk hydrogel wearers with silicone hydrogels. *Optom Vis Sci* 2004; 81 (12s): 31
47. Bergenske P, Long B, Dillehay S *et al.* Long-term clinical results: 3 years of up to 30-night continuous wear of lotrafilcon A silicone hydrogel and daily wear of low-Dk/t hydrogel lenses. *Eye Contact Lens* 2007; 33;2: 74-80.
48. Chalmers RL, Dillehay S, Long B *et al.* Impact of previous extended and daily wear schedules on signs and symptoms with high Dk lotrafilcon A lenses. *Optom Vis Sci* 2005; 82;6: 549-54.
49. Ladage PM, Yamamoto K, Ren DH *et al.* Effects of rigid and soft contact lens daily wear on corneal epithelium, tear lactate dehydrogenase, and bacterial binding to exfoliated epithelial cells. *Ophthalmology* 2001; 108 (7): 1279-88
50. Ladage PM, Ren DH, Petroll WM *et al.* Effects of eyelid closure and disposable and silicone hydrogel extended contact lens wear on rabbit corneal epithelial proliferation. *Invest Ophthalmol Vis Sci* 2003; 44 (5): 1843-9
51. Ren DH, Petroll WM, Jester JV *et al.* The relationship between contact lens oxygen permeability and binding of *Pseudomonas aeruginosa* to human corneal epithelial cells after overnight and extended wear. *CLAO J* 1999; 25;2: 80-100.
52. Ladage PM, Yamamoto K, Ren DH *et al.* Effects of rigid and soft contact lens daily wear on corneal epithelium, tear lactate dehydrogenase, and bacterial binding to exfoliated epithelial cells. *Ophthalmology* 2001; 108;7: 1279-88.

53. Ren D, Yamamoto K, Ladage P et al. Adaptive effects of 30-day extended wear of new hyper-O₂ transmissible RGP and soft contact lenses on bacterial binding and corneal epithelium. *Invest Ophthalmol Vis Sci* 2001; 42;4: s595.
54. Cavanagh HD, Ladage PM, Li SL et al. Effects of daily and overnight wear of a novel hyper oxygen-transmissible soft contact lens on bacterial binding and corneal epithelium: a 13-month clinical trial. *Ophthalmology* 2002; 109;11: 1957-69.
55. Ren DH, Yamamoto K, Ladage PM et al. Adaptive effects of 30-night wear of hyper-O₂ transmissible contact lenses on bacterial binding and corneal epithelium : A 1-year clinical trial. *Ophthalmology* 2002; 109;1: 27-39.
56. Cavanagh HD, Ladage P, Yamamoto K et al. Effects of daily and overnight wear of hyper-oxygen transmissible rigid and silicone hydrogel lenses on bacterial binding to the corneal epithelium: 13-month clinical trials. *Eye Contact Lens* 2003; 29;1 Suppl: S14-16; discussion S26-29, S192-194.
57. Holden BA, Williams L, Zantos SG: The etiology of transient endothelial changes in the human cornea. *Invest Ophthalmol Vis Sci* 1985; 26;10: 1354-9.
58. Ohya S, Nishimaki K, Nakayasu K, Kanai A: Non-contact specular microscopic observation for early response of corneal endothelium after contact lens wear. *Clao J* 1996; 22;2: 122-6.
59. Brennan NA. Endothelial bleb response to silicone hydrogels. Presentation at BCLA Conference 2008
60. Brennan NA, Coles ML, Ang JH: An evaluation of silicone-hydrogel lenses worn on a daily wear basis. *Clin Exp Optom* 2006; 89;1: 18-25.
61. Fonn D: Preventing contact lens dropouts. *Contact Lens Spectrum* 2002; 17;8: 26-32.
62. Fonn D, Dumbleton K: Dryness and discomfort with silicone hydrogel contact lenses. *Eye Contact Lens* 2003; 29;1 Suppl: S101-4; discussion S115-8, S192-4.
63. Fonn D: Targeting contact lens induced dryness and discomfort: what properties will make lenses more comfortable. *Optom Vis Sci* 2007; 84;4: 279-85.
64. Fonn D, Pritchard N, Brazeau D, Michaud L: Discontinuation of contact lens wear: The numbers, reasons and patient profiles. *Invest Ophthalmol Vis Sci* 1995; 36;4: S312.
65. Pritchard N, Fonn D, Brazeau D: Discontinuation of contact lens wear: a survey. *Int Cont Lens Clin* 1999; 26; 157 - 162.
66. Young G, Riley CM, Chalmers RL, Hunt C: Hydrogel lens comfort in challenging environments and the effect of refitting with silicone hydrogel lenses. *Optom Vis Sci* 2007; 84;4: 302-8.
67. Chalmers RL, Begley CG: Dryness symptoms among an unselected clinical population with and without contact lens wear. *Cont Lens Anterior Eye* 2006; 29;1: 25-30.
68. Richdale K, Sinnott LT, Skadahl E, Nichols JJ: Frequency of and factors associated with contact lens dissatisfaction and discontinuation. *Cornea* 2007; 26;2: 168-74.
69. Fonn D, Pritchard N, *et al.* Factors affecting the success of silicone hydrogels. In: Sweeney DF, ed. *Silicone hydrogels; the rebirth of extended wear contact lenses.* Oxford, UK: Butterworth-Heinemann 2000; 214-34

70. Ousler GW 3rd, Anderson RT, Osborn KE. The effect of senofilcon A contact lenses compared to habitual contact lenses on ocular discomfort during exposure to a controlled adverse environment. *Curr Med Res Opin.* 2008; 24 (2): 335-41
71. Schafer J, Mitchell GL, Chalmers RL et al. The stability of dryness symptoms after refitting with silicone hydrogel contact lenses over 3 years. *Eye Contact Lens.* 2007; 33 (5): 247-52
72. Malet F, Pagot R, Peyre C et al. Subjective experience with high-oxygen and low-oxygen permeable soft contact lenses in France. *Eye Contact Lens* 2003; 29;1: 55-9.
73. Guillon M, Maissa C: Use of silicone hydrogel material for daily wear. *Cont Lens Anterior Eye* 2007; 30;1: 5-10
74. Hall B, Jones S, Young G, Coleman S: The on-eye dehydration of proclear compatibles lenses. *CLAO J* 1999; 25;4: 233-7.
75. Lemp MA, Caffery B, Lebow K et al. Omaficon A (Proclear) soft contact lenses in a dry eye population. *CLAO J* 1999; 25;1: 40-7.
76. Fonn D, Situ P, Simpson T: Hydrogel lens dehydration and subjective comfort and dryness ratings in symptomatic and asymptomatic contact lens wearers. *Optom Vis Sci* 1999; 76;10: 700 - 704.
77. Chalmers RL, Dillehay S, Long B et al. Impact of previous extended and daily wear schedules on signs and symptoms with high Dk lotrafilcon A lenses. *Optom Vis Sci.* 2005; 82 (6): 549-54
78. Jones L, May C, Nazar L, Simpson T. In vitro evaluation of the dehydration characteristics of silicone hydrogel and conventional hydrogel contact lens materials. *Cont Lens Anterior Eye.* 2002; 25 (3): 147-56.
79. Morgan PB, Efron N. In vivo dehydration of silicone hydrogel contact lenses. *Eye Contact Lens* 2003; 29 (3): 173-6
80. Gonzalez-Meijome JM, Lopez-Aleman A, Almeida JB, Parafita MA, Refojo MF: Qualitative and quantitative characterization of the in vitro dehydration process of hydrogel contact lenses. *J Biomed Mater Res B Appl Biomater* 2007; 83;2: 512-26.
81. Dillehay SM: Does the level of available oxygen impact comfort in contact lens wear?: A review of the literature. *Eye Contact Lens* 2007; 33;3: 148-55.
82. Bruinsma GM, van der Mei HC, Busscher HJ: Bacterial adhesion to surface hydrophilic and hydrophobic contact lenses. *Biomaterials* 2001; 22;24: 3217-24.
83. Court JL, Redman RP, Wang JH et al. A novel phosphorylcholine-coated contact lens for extended wear use. *Biomaterials* 2001; 22;24: 3261-72.
84. Cheng L, Muller SJ, Radke CJ: Wettability of silicone-hydrogel contact lenses in the presence of tear-film components. *Curr Eye Res* 2004; 28;2: 93-108.
85. Lorentz H, Rogers R, Jones L: In vitro deposition of lipid onto contact lens materials can lower contact angle wettability of surface-modified silicone hydrogel contact lens materials. *Invest Ophthalmol Vis Sci* 2006; 46;E-Abstract 2389.
86. Lorentz H, Rogers R, Jones L: The impact of lipid on contact angle wettability. *Optom Vis Sci* 2007; 84;10: 946-53.
87. Maldonado-Codina C, Morgan PB: In vitro water wettability of silicone hydrogel contact lenses

determined using the sessile drop and captive bubble techniques. *J Biomed Mater Res A* 2007; 83;2: 496-502.

88. Huth S, Wagner H: Identification and removal of deposits on polydimethylsiloxane silicone elastomer lenses. *Int Contact Lens Clin* 1981; 8;7/8: 19-26.

89. Jones L, Senchyna M, Glasier MA et al. Lysozyme and lipid deposition on silicone hydrogel contact lens materials. *Eye Contact Lens* 2003; 29;1 Suppl: S75-S79.

90. Carney FP, Nash WL, Sentell KB: The adsorption of major tear film lipids in vitro to various silicone hydrogels over time. *Invest Ophthalmol Vis Sci* 2008; 49;1: 120-4.

91. Rae ST, Huff JW: Studies on initiation of silicone elastomer lens adhesion in vitro: binding before the indentation ring. *CLAO J* 1991; 17;3: 181 - 186.

92. Jones L, Subbaraman LN, Rogers R, Dumbleton K: Surface treatment, wetting and modulus of silicone hydrogels. *Optician* 2006; 232;6067: 28 - 34.

93. Valint PL, Jr., Grobe GL, 3rd, Ammon DM, Jr., Moorehead M. Plasma surface treatment of silicone hydrogel contact lenses. 2001; US Patent # 6,193,369.

94. Grobe G, Kunzler J, Seelye D, Salamone J: Silicone hydrogels for contact lens applications. *Polymeric Materials Science and Engineering* 1999; 80; 108 - 109.

95. Lopez-Aleman A, Compan V, Refojo MF: Porous structure of Purevision versus Focus Night & Day and conventional hydrogel contact lenses. *J Biomed Mater Res (Appl Biomater)* 2002; 63; 319 - 325.

96. Gonzalez-Meijome JM, Lopez-Aleman A, Almeida JB, Parafita MA, Refojo MF: Microscopic observation of unworn siloxane-hydrogel soft contact lenses by atomic force microscopy. *J Biomed Mater Res B Appl Biomater* 2006; 76;2: 412-8.

97. Nicolson PC: Continuous wear contact lens surface chemistry and wearability. *Eye Contact Lens* 2003; 29;1 Suppl: S30-2; discussion S57-9, S192-4.

98. Jones L: A new silicone hydrogel lens comes to market. *Contact Lens Spectrum* 2007; 22;10: 23.

99. Steffen R, Schnider C: A next generation silicone hydrogel lens for daily wear. Part 1 - Material properties. *Optician* 2004; 227;5954: 23 - 25.

100. Osborn K & Veys J. A new silicone hydrogel lens for contact lens related dryness. Material properties. *OPTICIAN* 2005;229:6004 39-41

101. Jones L: Comfilcon A: a new silicone hydrogel material. *Contact Lens Spectrum* 2007; 22;8: 21.

102. French K: Contact lens material properties part 1: Wettability. *Optician* 2005; 230;6022: 20 - 28.

103. Jones L, Senchyna M: Soft contact lens solutions review: Part 1 - components of modern care regimens. *Optometry in Practice* 2007; 8; 45 - 56.

104. Jones L, Senchyna M: Soft contact lens solutions review: Part 2 - modern generation care systems. *Optometry in Practice* 2008; 9; 43 - 62.

105. Jones L: Understanding the link between wettability and lens comfort. *Contact Lens Spectrum* 2007; 22;6 (supp): s4-s6.

106. Ross G, *et al.* Silicone Hydrogels: Trends in Products and Properties. Poster presented at BCLA 29th Clinical Conference & Exhibition, Brighton, UK; 3-5 June, 2005
107. Walsh JE, Koehler LV, Fleming DP, Bergmanson JP: Novel method for determining hydrogel and silicone hydrogel contact lens transmission curves and their spatially specific ultraviolet radiation protection factors. *Eye Contact Lens* 2007; 33;2: 58-64.
108. Moore L, Ferreira JT: Ultraviolet (UV) transmittance characteristics of daily disposable and silicone hydrogel contact lenses. *Cont Lens Anterior Eye* 2006; 29;3: 115-22.
109. Walsh JE, Bergmanson JP, Saldana G, Jr., Gaume A: Can UV radiation-blocking soft contact lenses attenuate UV radiation to safe levels during summer months in the southern United States? *Eye Contact Lens* 2003; 29;1 Suppl: S174-9; discussion S190-1, S192-4.
110. Quesnel NM, Fares F, Verret E, Giasson C: Evaluation of the spectral transmittance of UV-absorbing disposable contact lenses. *Clao J* 2001; 27;1: 23-9.
111. Harris MG, Chin RS, Lee DS, Tam MH, Dobkin CE: Ultraviolet transmittance of the Vistakon disposable contact lenses. *Cont Lens Anterior Eye* 2000; 23;1: 10-5.
112. Harris MG, Haririfar M, Hirano KY: Transmittance of tinted and UV-blocking disposable contact lenses. *Optom Vis Sci* 1999; 76;3: 177-80.
113. Bergmanson JP, Sheldon TM: Ultraviolet radiation revisited. *Clao J* 1997; 23;3: 196-204.
114. Kwok LS, Daszynski DC, Kuznetsov VA, Pham T, Ho A, Coroneo MT: Peripheral light focusing as a potential mechanism for phakic dysphotopsia and lens phototoxicity. *Ophthalmic Physiol Opt* 2004; 24;2: 119-29.
115. Coroneo MT, Muller-Stolzenburg NW, Ho A: Peripheral light focusing by the anterior eye and the ophthalmohelioses. *Ophthalmic Surg* 1991; 22;12: 705-11.
116. Kwok LS, Coroneo MT: A model for pterygium formation. *Cornea* 1994; 13;3: 219-24.
117. Kwok LS, Kuznetsov VA, Ho A, Coroneo MT: Prevention of the adverse photic effects of peripheral light-focusing using UV-blocking contact lenses. *Invest Ophthalmol Vis Sci* 2003; 44;4: 1501-7.
118. ANSI/Z80.3 – 1996 (R1999) Non-prescription sunglasses and fashion eyewear – requirements
119. ISO 8599:1994 Optics and optical instruments – contact lenses – Determination of the spectral and luminous transmittance